

# Additional documentation for the Gay-Berne ellipsoidal potential as implemented in LAMMPS

Mike Brown, Sandia National Labs, April 2007

The Gay-Berne anisotropic LJ interaction between pairs of dissimilar ellipsoidal particles is given by

$$U(\mathbf{A}_1, \mathbf{A}_2, \mathbf{r}_{12}) = \mathbf{U}_r(\mathbf{A}_1, \mathbf{A}_2, \mathbf{r}_{12}, \gamma) \cdot \eta_{12}(\mathbf{A}_1, \mathbf{A}_2, v) \cdot \chi_{12}(\mathbf{A}_1, \mathbf{A}_2, \mathbf{r}_{12}, \mu)$$

where  $\mathbf{A}_1$  and  $\mathbf{A}_2$  are the transformation matrices from the simulation box frame to the body frame and  $\mathbf{r}_{12}$  is the center to center vector between the particles.  $U_r$  controls the shifted distance dependent interaction based on the distance of closest approach of the two particles ( $h_{12}$ ) and the user-specified shift parameter gamma:

$$U_r = 4\epsilon(\varrho^{12} - \varrho^6)$$

$$\varrho = \frac{\sigma}{h_{12} + \gamma\sigma}$$

Let the shape matrices  $\mathbf{S}_i = \text{diag}(\mathbf{a}_i, \mathbf{b}_i, \mathbf{c}_i)$  be given by the ellipsoid radii. The  $\eta$  orientation-dependent energy based on the user-specified exponent  $v$  is given by

$$\eta_{12} = \left[ \frac{2s_1 s_2}{\det(\mathbf{G}_{12})} \right]^{v/2},$$

$$s_i = [a_i b_i + c_i c_i][a_i b_i]^{1/2},$$

and

$$\mathbf{G}_{12} = \mathbf{A}_1^T \mathbf{S}_1^2 \mathbf{A}_1 + \mathbf{A}_2^T \mathbf{S}_2^2 \mathbf{A}_2 = \mathbf{G}_1 + \mathbf{G}_2.$$

Let the relative energy matrices  $\mathbf{E}_i = \text{diag}(\epsilon_{ia}, \epsilon_{ib}, \epsilon_{ic})$  be given by the relative well depths (dimensionless energy scales inversely proportional to the well-depths of the respective orthogonal configurations of the interacting

molecules). The  $\chi$  orientation-dependent energy based on the user-specified exponent  $\mu$  is given by

$$\chi_{12} = [2\hat{\mathbf{r}}_{12}^T \mathbf{B}_{12}^{-1} \hat{\mathbf{r}}_{12}]^\mu,$$

$$\hat{\mathbf{r}}_{12} = \mathbf{r}_{12}/|\mathbf{r}_{12}|,$$

and

$$\mathbf{B}_{12} = \mathbf{A}_1^T \mathbf{E}_1^2 \mathbf{A}_1 + \mathbf{A}_2^T \mathbf{E}_2^2 \mathbf{A}_2 = \mathbf{B}_1 + \mathbf{B}_2.$$

Here, we use the distance of closest approach approximation given by the Perram reference, namely

$$h_{12} = r - \sigma_{12}(\mathbf{A}_1, \mathbf{A}_2, \mathbf{r}_{12}),$$

$$r = |\mathbf{r}_{12}|,$$

and

$$\sigma_{12} = [\frac{1}{2}\hat{\mathbf{r}}_{12}^T \mathbf{G}_{12}^{-1} \hat{\mathbf{r}}_{12}]^{-1/2}$$

Forces and Torques: Because the analytic forces and torques have not been published for this potential, we list them here:

$$\mathbf{f} = -\eta_{12}(\mathbf{U}_r \cdot \frac{\partial \chi_{12}}{\partial \mathbf{r}} + \chi_{12} \cdot \frac{\partial \mathbf{U}_r}{\partial \mathbf{r}})$$

where the derivative of  $U_r$  is given by (see Allen reference)

$$\frac{\partial U_r}{\partial r} = \frac{\partial U_{SLJ}}{\partial r} \hat{\mathbf{r}}_{12} + r^{-2} \frac{\partial U_{SLJ}}{\partial \varphi} [\kappa - (\kappa^T \cdot \hat{\mathbf{r}}_{12}) \hat{\mathbf{r}}_{12}],$$

$$\frac{\partial U_{SLJ}}{\partial \varphi} = 24\epsilon(2\varrho^{13} - \varrho^7)\sigma_{12}^3/2\sigma,$$

$$\frac{\partial U_{SLJ}}{\partial r} = 24\epsilon(2\varrho^{13} - \varrho^7)/\sigma,$$

and

$$\kappa = \mathbf{G}_{12}^{-1} \cdot \mathbf{r}_{12}.$$

The derivate of the  $\chi$  term is given by

$$\frac{\partial \chi_{12}}{\partial r} = -r^{-2} \cdot 4.0 \cdot [\iota - (\iota^T \cdot \hat{\mathbf{r}}_{12}) \hat{\mathbf{r}}_{12}] \cdot \mu \cdot \chi_{12}^{(\mu-1)/\mu},$$

and

$$\iota = \mathbf{B}_{12}^{-1} \cdot \mathbf{r}_{12}.$$

The torque is given by:

$$\begin{aligned}\tau_i &= \mathbf{U}_r \eta_{12} \frac{\partial \chi_{12}}{\partial \mathbf{q}_i} + \chi_{12} (\mathbf{U}_r \frac{\partial \eta_{12}}{\partial \mathbf{q}_i} + \eta_{12} \frac{\partial \mathbf{U}_r}{\partial \mathbf{q}_i}), \\ \frac{\partial U_r}{\partial \mathbf{q}_i} &= \mathbf{A}_i \cdot (-\kappa^T \cdot \mathbf{G}_i \times \mathbf{f}_k), \\ \mathbf{f}_k &= -\mathbf{r}^{-2} \frac{\delta \mathbf{U}_{SLJ}}{\delta \varphi} \kappa,\end{aligned}$$

and

$$\frac{\partial \chi_{12}}{\partial \mathbf{q}_i} = 4.0 \cdot r^{-2} \cdot \mathbf{A}_i (-\iota^T \cdot \mathbf{B}_i \times \iota).$$

For the derivative of the  $\eta$  term, we were unable to find a matrix expression due to the determinant. Let  $a_{mi}$  be the mth row of the rotation matrix  $A_i$ . Then,

$$\frac{\partial \eta_{12}}{\partial \mathbf{q}_i} = \mathbf{A}_i \cdot \sum_m \mathbf{a}_{mi} \times \frac{\partial \eta_{12}}{\partial \mathbf{a}_{mi}} = \mathbf{A}_i \cdot \sum_m \mathbf{a}_{mi} \times \mathbf{d}_{mi},$$

where  $d_{mi}$  represents the mth row of a derivative matrix  $D_i$ ,

$$\mathbf{D}_i = -\frac{1}{2} \cdot \left( \frac{2s1s2}{\det(\mathbf{G}_{12})} \right)^{v/2} \cdot \frac{v}{\det(\mathbf{G}_{12})} \cdot \mathbf{E},$$

where the matrix  $E$  gives the derivate with respect to the rotation matrix,

$$\mathbf{E} = [\mathbf{e}_{my}] = \frac{\partial \eta_{12}}{\partial \mathbf{A}_i},$$

and

$$e_{my} = \det(\mathbf{G}_{12}) \cdot \text{trace}[\mathbf{G}_{12}^{-1} \cdot (\hat{\mathbf{p}}_y \otimes \mathbf{a}_m + \mathbf{a}_m \otimes \hat{\mathbf{p}}_y) \cdot \mathbf{s}_{mm}^2].$$

Here,  $p_v$  is the unit vector for the axes in the lab frame ( $p1 = [1, 0, 0]$ ,  $p2 = [0, 1, 0]$ , and  $p3 = [0, 0, 1]$ ) and  $s_{mm}$  gives the  $m$ th radius of the ellipsoid  $i$ .